

Environmental Science at Beamline 10.3.2

M. Marcus, A. A. Macdowell, R. Sublett, R. Celestre, E. Domning,
T. Miller, K. Franck, H. A. Padmore
Advanced Light Source
A. Manceau
LGIT, Université Joseph Fourier


*Themes: How are toxic elements sequestered in the environment?
How do living organisms deal with toxics?
What happens when a toxic site is remediated?*

*What we get from 10.3.2:
XRF mapping: Where is it? What other elements is it with?
EXAFS: What is its chemical form? What valence? What's it attached to?
XRD (new): What mineral is it with?*


Unlocking Metal Sequestration in Soils

A. Manceau, N. Tamura, M.A. Marcus, A.A. MacDowell
G. Sposito, and H.A. Padmore

How are trace elements naturally sequestered in soils?
Ferromanganese nodules are common sinks.



Photomicrograph of a soil thin section containing ferromanganese nodules.



Typical back-scatter electron (BSE) images of soil micronodules.

Objectives

To determine quantitatively the forms of Zn and Ni in typical soil micronodules.


Ultimately, to compare sequestration mechanisms of trace metals from a variety of continental soils.

Approach


Synergistic use of three non-invasive synchrotron-based techniques:

- X-ray microfluorescence (SRXF)
- X-ray microdiffraction (XRD and SXRD)
- X-ray absorption spectroscopy (EXAFS and EXAFS)


Elemental Maps (XRF)




Mineral Species Maps (XRD)



Matrix minerals



Sequestered forms of metals (EXAFS)



Conclude: In soil nodules, Ni occurs as 1 species (Ni-lithiophorite), Zn as several.